



DATA SERVICES ARTICLE

GESLA Version 3: A major update to the global higher-frequency sea-level dataset

Ivan D. Haigh¹ | Marta Marcos² | Stefan A. Talke³ | Philip L. Woodworth⁴ |
John R. Hunter⁵ | Ben S. Hague⁶ | Arne Arns⁷ | Elizabeth Bradshaw⁸ |
Philip Thompson⁹

¹School of Ocean and Earth Science, University of Southampton, National Oceanography Centre, Southampton, UK

²IMEDEA (UIB-CSIC), Esporles, Balearic Islands, Spain

³Department of Civil and Environmental Engineering, California Polytechnic State University, San Luis Obispo, California, USA

⁴National Oceanography Centre, Liverpool, UK

⁵Institute for Marine and Antarctic Studies, University of Tasmania, Hobart, Tasmania, Australia

⁶Australian Bureau of Meteorology, Melbourne, Victoria, Australia

⁷Faculty of Agricultural and Environmental Sciences, University of Rostock, Rostock, Germany

⁸British Oceanographic Data Centre, National Oceanography Centre, Liverpool, UK

⁹Department of Oceanography, University of Hawai'i at Mānoa, Honolulu, Hawai'i, USA

Correspondence

Ivan D. Haigh, School of Ocean and Earth Science, University of Southampton, National Oceanography Centre, European Way, Southampton SO14 3ZH, UK.

Email: i.d.haigh@soton.ac.uk

Abstract

This paper describes a major update to the quasi-global, higher-frequency sea-level dataset known as GESLA (Global Extreme Sea Level Analysis). Versions 1 (released 2009) and 2 (released 2016) of the dataset have been used in many published studies, across a wide range of oceanographic and coastal engineering-related investigations concerned with evaluating tides, storm surges, extreme

DATASET DETAILS

This dataset is called GESLA (Global Extreme Sea Level Analysis), and it contains high-frequency (at least hourly) sea-level information from tide gauge stations distributed worldwide. This update, Version 3, contains 91,021 years of data from 5,119 records, a significant increase from Version 2. The dataset is available from <https://www.gesla.org> and is also distributed by the British Oceanographic Data Centre (BODC) with Digital Object Identifiers (DOI) as follows:

Identifiers: DOIs:

Part 1, [10.5285/d21a496a-a48e-1f21-e053-6c86abc08512](https://doi.org/10.5285/d21a496a-a48e-1f21-e053-6c86abc08512);

Part 2, [10.5285/d21a496a-a48f-1f21-e053-6c86abc08512](https://doi.org/10.5285/d21a496a-a48f-1f21-e053-6c86abc08512)

Websites:

Part 1, https://www.bodc.ac.uk/data/published_data_library/catalogue/10.5285/d21a496a-a48e-1f21-e053-6c86abc08512/;

Part 2, https://www.bodc.ac.uk/data/published_data_library/catalogue/10.5285/d21a496a-a48f-1f21-e053-6c86abc08512/

Title: GESLA (Global Extreme Sea Level Analysis) higher-frequency sea-level dataset, Version 3.

Publisher: NERC EDS British Oceanographic Data Centre NOC

Publication year: 2021

Resource type: Dataset

Version: 3

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2022 The Authors. *Geoscience Data Journal* published by Royal Meteorological Society and John Wiley & Sons Ltd.

Funding information

National Oceanic and Atmospheric Administration Global Ocean Monitoring and Observation Program, Grant/Award Number: NA11NMF4320128; European Regional Development Fund and Ministerio de Ciencia, Innovación y Universidades – Agencia Estatal de Investigación through the MOCCA project, Grant/Award Number: RTI2018-093941-B-C31; National Science Foundation, Grant/Award Number: 2013280 and 1455350; NERC-funded CHANCE Project, Grant/Award Number: NE/S010262/1

sea levels, and other related processes. The third version of the dataset (released 2021), presented here, contains double the number of years of data, and nearly four times the number of records, compared to Version 2. The dataset consists of records obtained from multiple sources around the world. This paper describes the assembly of the dataset, its processing, and its format, and outlines potential future improvements.

KEYWORDS

sea level records, sea level rise, storm surges, storm tides, tide gauge

1 | INTRODUCTION

Having access to high-quality sea-level measurements worldwide is vital for many oceanographic and coastal applications. For example, sea-level records form the basis of our understanding of changes in mean sea level, which affects the livelihoods of hundreds of millions of people living in the world's coastal regions and is one of the key indicators of climate change (Oppenheimer et al., 2019). Coastal sea-level extremes are among the costliest and potentially most hazardous impacts affecting densely populated coastal regions (Wong et al., 2014). Analyses of sea-level records help engineers and coastal managers define flood defence heights and other coastal protection measures. Measurements of sea level are used to map the timing and heights of astronomical tides and calibrate and validate both operational and scientific numerical models of oceanic processes (Muis et al., 2020). Furthermore, coastal sea-level measurements form a key component of the data used in nautical charts and geodetic surveys, and influence legal definitions of shoreline boundaries (Shalowitz, 1962). Building on an earlier study (Woodworth et al., 2017), this paper is concerned with extending a global dataset of higher-frequency (at least hourly) sea-level records from tide gauges at as many locations as possible worldwide.

The international body responsible for coordinating collection and access to in situ sea-level records is the Global Sea Level Observing System (GLOSS), which was established by the UNESCO Intergovernmental Oceanographic Commission (IOC) in 1985 to support a broad research and operational user base. Multiple GLOSS data centres contribute to the aggregation of global sea-level datasets with varying temporal resolutions and levels of quality control. Global datasets of monthly and annual mean sea levels have been available for many decades via the Permanent Service for Mean Sea Level (PSMSL). Established in 1933, PSMSL has been responsible for the

collection of mean sea-level data from global tide gauges (Holgate et al., 2013) and has been used, with altimeter records, in most past mean sea-level trend and variability studies. PSMSL has always had good coverage globally because, historically, tide gauge operators have been more willing to share monthly mean data, rather than higher-frequency data. However, higher-frequency data are required for the study of ocean tides, storm surges, and extreme sea levels (Woodworth et al., 2019). The GLOSS dataset for research-quality hourly sea-level data are the Joint Archive for Sea Level (Caldwell et al., 2015), which was established in 1987 and is hosted by the University of Hawaii Sea Level Center (UHSLC). This dataset is composed of nearly 18,000 years of hourly sea-level data from 696 records in 97 countries. These data have been inspected for outliers, timing issues, and datum shifts, and efforts have been made to reconcile quality issues with the data originators. The locations of records in the UHSLC dataset are distributed globally, with care given to balance global coverage with the time-intensive process of quality assessment. Thus, the UHSLC dataset excludes many records in densely sampled regions to provide global coverage while maintaining an update cycle of approximately 2 years.

The GESLA (Global Extreme Sea Level Analysis) project was established, over a decade ago, to increase access to a greater volume of the global hourly and even higher-frequency sea-level data, than is available in the UHSLC dataset. The original aim of the project was to assemble as many higher-frequency sea-level records as were readily available into a common format with consistent quality control flags, to make it easier for researchers to maximize the geographic density of data, capturing extreme sea levels on a global scale. The first GESLA dataset, denoted GESLA-1, was assembled in 2009 and contained 21,197 years of higher-frequency measurements from 675 records. The majority of the data were obtained by ingesting UHSLC and other GLOSS data. The GLOSS

datasets were then supplemented by a small number of other records obtained from national data centres or from contributions received from colleagues in the sea-level community. GESLA-1 was first used in a study of sea-level extremes by Menendez and Woodworth (2010). Subsequent publications based on GESLA-1 included, for example, Hunter et al. (2013), Mawdsley et al. (2015) and Marcos et al. (2015), and GESLA-1 was used in the Intergovernmental Panel on Climate Change's (IPCC) Fifth Assessment Report (Church et al., 2013; Rhein et al., 2013; Wong et al., 2014).

After some years, it became apparent that GESLA-1 needed updating to include additional data and to extend its coverage in under-represented areas. Thus GESLA-2 was assembled in 2015 and 2016. The compilation of GESLA-2 is described in detail in Woodworth et al. (2017). This second version contained almost twice the amount of data compared to the first. GESLA-2 contained 39,151 years of higher-frequency measurements of sea level from 1,355 records; again, the UHSLC dataset made up a significant proportion of this database. Since its release in early 2016, GESLA-2 has been used in a wide range of ocean research, examples including:

1. Assessment of temporal and spatial changes in extreme sea levels and links to regional climate (e.g., Marcos & Woodworth, 2017; Rashid et al., 2021).
2. Calculation of extreme sea-level return periods and sea-level allowances (e.g., Tsitsikas, 2018; Wahl et al., 2017; Woodworth et al., 2021).
3. Provision of information for flood inundation studies (e.g., Hunter et al., 2017).
4. Analysis of nonlinear interactions between tides and nontidal residuals or skew surges (e.g., Arns et al., 2020; Santamaria-Aguilar & Vafeidis, 2019).
5. Investigations of changes in ocean tidal constituents and levels (e.g., Ray, 2020; Schindelegger et al., 2018).
6. Examinations of the magnitude and changes in the perigean and nodal inter-annual tidal cycles (e.g., Peng et al., 2019; Woodworth & Hibbert, 2018).
7. Validation of regional and global ocean tide and tide/surge hydrodynamic models (e.g., Muis et al., 2020; Piccioni et al., 2018).
8. Assessment of compound flooding from coastal, fluvial, and pluvial sources (e.g., Ward et al., 2018).
9. Other applications (e.g., Tadesse et al., 2020; Wolff et al., 2018).

GESLA-2 data has also been used in the IPCC Special Report on Ocean and the Cryosphere (Oppenheimer et al., 2019), and in the Sixth Assessment Report (Fox-Kemper et al., 2021). Furthermore, a secondary database of tidal constituents has been derived from GESLA-2 by

Piccioni et al. (2019) and another for skew surges has also been made available through the GESLA website, after Woodworth et al. (2017). All the studies that the authors are aware of that have used the GESLA dataset to date are listed on <https://www.gesla.org>. In 2016, GESLA was made an official GLOSS dataset.

In this paper, we describe the development of Version 3 of the dataset, which provides a major update. Section 2 of this paper describes the data sources, the data processing, and the revised GESLA data format. Access to the data set is described in Section 3. A discussion and conclusions are given in Section 4.

2 | DATA DESCRIPTION AND DEVELOPMENT

Here, we describe the data sources, record locations, and the number of years of data (Section 2.1), we outline the data processing and format (Section 2.2), we describe the usage licences (Section 2.3), and we discuss the dataset in regards to the recently established FAIR (findable, accessible, interoperable, and reusable) data principles (Section 2.4).

2.1 | Data sources

We obtained the higher-frequency sea-level dataset for GESLA-3 from 36 international and national data providers (Table 1). Providers are ordered by the number of years of sea-level data available (see Table 2). Below, we use the abbreviated names of the providers; readers should refer to Table 1 for their full names. We define the length of a sea-level dataset for a particular record, as being the number of years available; a year is a calendar year containing one or more sea-level measurements for that particular record. We use the term record to refer to a sea-level dataset at a particular tide gauge. A specific tide gauge station can have more than one record; either because (a) a duplicate record for that station is available from different providers or (b) because sometimes sea-level time series for the same station is split into different records when there are datum jumps or changes in the location or instrument (i.e., the UHSLC dataset contains such records, and these are denoted by letters, A, B, C, etc. after the station code).

Data were obtained and processed as follows. First, full records were downloaded again from all the sources used to compile GESLA-2 (Table 2 in Woodworth et al., 2017), except where noted below. Therefore, any changes to quality control or datums made since 2015/16 are reflected in GESLA-3. GESLA-2 included 191 records from the GLOSS Delayed Model dataset (source

TABLE 1 Information on data providers, licences, and data use. Grey shading indicates new data sources not in GESLA-2

Number	Abbreviated name	Full name	Website	Country	Download method	Licence	Use
1	UHSLC	University of Hawaii Sea level Center	https://uhslc.soest.hawaii.edu	Global	Downloaded each record automatically via API (ERDDAP server)	Specified on website: https://uhslc.soest.hawaii.edu/erddap/global_daily_rqds.html	Research and consultancy
2	NOAA	National Oceanic and Atmospheric Administration	https://api.tidesandcurrents.noaa.gov/api/prod/	United States of America	Downloaded each record automatically via API	Unspecified	Research and consultancy
3	CMEMS	Copernicus Marine Environment Monitoring Service	https://resources.marine.copernicus.eu/?option=com_csw&view=details&product_id=INSITU_GLO_NRT_OBSERVATIONS_013_030	Europe	Download netcdf files from ftp site	Specified in netcdf data files	Research (for consultancy contact data owners directly)
4	MEDS	Marine Environmental Data Section	https://isdm-gdsi.gc.ca/isdm-gdsi/twl-mne/inventory-inventaire/index-eng.htm	Canada	Downloaded each record manually from website (in 10-year blocks)	Specified on website: https://www.qc.dfo-mpo.gc.ca/tides/en/licence-agreement	Research and consultancy
5	USGS	United States Geological Survey	http://waterdata.usgs.gov/nwis/uv	United States of America	Downloaded each record manually from website	Unspecified	Research and consultancy
6	BOM	Bureau of Meteorology	http://www.bom.gov.au/oceanography/projects/absimp/absimp.shtml	Australia and Pacific Islands	Obtained directly from BOM	Unspecified	Research and consultancy
7	RWS	Rijkswaterstaat	https://opendap.delta.res.nl/thredds/catalog/opendap/rijkswaterstaat/waterbase/27_Waterhoogte_in_cm_to.v_normaal_amsterdams_pegel_in_oppevlaktewater/nc/catalog.html	The Netherlands	Downloaded each record manually from website	Unspecified	Research and consultancy
8	JODC_JMA	Japan Oceanographic Data Center, Japan Meteorological Agency	https://jdoss1.jodc.go.jp/vpage/tide.html	Japan	Downloaded each site manually from website (in 10-year blocks)	Specified on website: https://jdoss1.jodc.go.jp/vpage/tide.html	Research and consultancy

TABLE 1 (Continued)

Number	Abbreviated name	Full name	Website	Country	Download method	Licence	Use
9	SMHI	Swedish Meteorological and Hydrological Institute	https://www.smhi.se/data/oceanografi/ladda-ner-oceanografiska-observationer/#param=scalelevelMI&nutes,stations=all	Sweden	Manually downloaded each record from website	http://www.smhi.se/data/oppna-data/information-om-oppna-data/villkor-for-anvandning-1.30622	Research and consultancy
10	REFMAR	Réseaux de référence des observations marégraphiques (Reference networks for tidal observations)	http://refmar.shom.fr/en	France	Downloaded each record manually from website	Unspecified	Research and consultancy
11	BODC	British Oceanographic Data Centre	https://www.bodc.ac.uk/data/hosted_data_systems/sea_level/uk_tide_gauge_network/	United Kingdom of Great Britain and Northern Ireland	Downloaded each record manually from website	https://www.nationalarchives.gov.uk/doc/open-government-licence/version/2/	Research and consultancy
12	CDWR	California Department of Water Resources	https://cdec.water.ca.gov/	United States of America	Downloaded each record manually from website	Unspecified	Research and consultancy
13	JODC_JCG	Japan Oceanographic Data Center, Japan Coast Guard	https://jdoss1.jodc.go.jp/vpage/tide.html	Japan, Antarctica	Downloaded each record manually from website	Specified on website: https://jdoss1.jodc.go.jp/vpage/tide.html	Research and consultancy
14	NHS	Norwegian Hydrographic Service	http://api.sehavniva.no/tideapi_en.html	Norway	Downloaded each record automatically via API and combined with historic data obtained directly	https://creativecommons.org/licenses/by/4.0/deed.en	Research and consultancy
15	JODC_GIAJ	Japan Oceanographic Data Center, Geospatial Information Authority of Japan	https://jdoss1.jodc.go.jp/vpage/tide.html	Japan	Downloaded each record manually from website	Specified on website: https://jdoss1.jodc.go.jp/vpage/tide.html	Research and consultancy
16	WSV	Wasserstraßen- und Schifffahrtsverwaltung des Bundes (Federal Waterway and Shipping Administration)	https://www.kuestendaten.de/DE/dynamisch/Funktionen/Liste_der_vorhandenen_Daten/index.php.html	Germany	Downloaded each record manually from website	Unspecified	Research and consultancy

(Continues)

TABLE 1 (Continued)

Number	Abbreviated name	Full name	Website	Country	Download method	Licence	Use
17	JODC_PAHB	Japan Oceanographic Data Center, Ports and Harbours Bureau	https://jdoss1.jodc.go.jp/vpage/tide.html	Japan	Downloaded each record manually from website	Specified on website: https://jdoss1.jodc.go.jp/vpage/tide.html	Research and consultancy
18	SFWMD	South Florida Water Management District	https://www.sfwmd.gov/science-data/dbhydro	United States of America	Downloaded each record manually from website	Unspecified	Research and consultancy
19	ISPRA	Istituto Superiore per la Protezione e la Ricerca Ambientale (Higher Institute for Environmental Protection and Research)	https://mareografico.it	Italy	Downloaded each record manually from website	N/A	Research and consultancy
20	IEO	Instituto Español de Oceanografía (Spanish Institute of Oceanography)	https://www.seadatanet.org	Spain	Downloaded each record manually from website	Unspecified	Research and consultancy
21	DA	Data Archeology	N/A	United States of America, Spain, United Kingdom of Great Britain and Northern Ireland	Obtained directly from authors	N/A	Research and consultancy
22	UNAM	National Autonomous University of Mexico	http://www.mareografi.co.unam.mx/portal/	Mexico	Downloaded each record manually from website	Specified on website http://www.mareografico.unam.mx/portal/	Research and consultancy
23	FMI	Finnish Meteorological Institute	https://en.ilmatieteenlaitos.fi/download-observations	Finland	Downloaded each record manually from website (in 15-year blocks)	https://en.ilmatieteenlaitos.fi/open-data-licence	Research and consultancy
24	DMI	Danish Meteorological Institute	http://ocean.dmi.dk/english/index.php	Denmark	Data obtained directly	Unspecified	Research and consultancy

TABLE 1 (Continued)

Number	Abbreviated name	Full name	Website	Country	Download method	Licence	Use
25	BFG	Bundesanstalt Für Gewässerkunde (Federal Institute of Hydrology)	https://www.bafg.de/EN/03_The_%20BfG/the_bfg.html	Germany	Data obtained directly	Unspecified	Research and consultancy
26	MI_C	Marine Institute (Coastal sites)	https://erddap.marine.ie/erddap/tabledap/IrishNationalTideGaugeNetwork.html	Ireland	Downloaded each record automatically via API (ERDDAP server)	https://creativecommons.org/licenses/by/4.0/	Research and consultancy
27	CCO	Coastal Channel Observatory	https://coastalmonitoring.org.org/realtime/data/	United Kingdom of Great Britain and Northern Ireland	Downloaded each record manually from website	https://www.nationalarchives.gov.uk/doc/open-government-licence/version/2/	Research and consultancy
28	NOC	National Oceanography Centre	https://noc.ac.uk	United Kingdom of Great Britain and Northern Ireland, Egypt, Ukraine	Data obtained directly	Unspecified	Research and consultancy
29	NWFWMD	North West Florida Water Management Department	https://nwfwater.com/Data-Publications/Hydrologic-Data/Active-Stations-Map	United States of America	Downloaded each record manually from website	Unspecified	Research and consultancy
30	ESEAS	European Sea-Level Service	https://www.bodc.ac.uk/projects/data_management/european/eseas/	Poland, Turkey, Croatia	Copied from GESLA2	Unspecified	Research and consultancy
31	ICG	Icelandic Coast Guard Hydrographic and Maritime Safety Department	http://www.lhg.is/english/about-us/	Iceland	Data obtained directly	Unspecified	Research and consultancy
32	UZ	University of Zagreb	https://www.pmf.unizg.hr/geof/en#	Croatia	Data obtained directly	Unspecified	Research (for consultancy contact data owners directly)

(Continues)

TABLE 1 (Continued)

Number	Abbreviated name	Full name	Website	Country	Download method	Licence	Use
33	NCDEM	North Carolina Department of Emergency Management	https://www.ncdps.gov/nceem	United States of America	Downloaded each record manually from website	Unspecified	Research and consultancy
34	CV	City of Venice, Tide Forecasts and Reporting Center	https://www.comune.venezia.it/node/6214	Italy	Downloaded manually from website	https://creativecommons.org/licenses/by-nc-sa/3.0/it/	Research (for consultancy contact data owners directly)
35	MI_R	Marine Institute (River Sites)	https://erddap.marine.ie/erddap/tabledap/IrishNationalTideGaugeNetworkRiverGauges.html	Ireland	Automatically load in data using ERDDAP	https://creativecommons.org/licenses/by/4.0/	Research and consultancy
36	GLOSS	Global Sea Level Observing System	https://gloss-sealevel.org	Greenland	Copied from GESLA2	Unspecified	Research and consultancy

Note: Bureau of Meteorology represents the same provider as the National Tidal Centre Australia in Table 2 of Woodworth et al. (2017).

1 glossdm-bodc, see Table 2 in Woodworth et al., 2017). However, this dataset has not been updated for many years and data from all but two of these records (Aasiaat and Maniitsoq in Greenland) are now available from other sources (see Table 1). Hence, we only included these two records in GESLA-3. GESLA-2 also included two datasets for Australia (source 28 johnhunter and 29 national_tidal_centre, see Table 2 in Woodworth et al., 2017). We did not include either of these in GESLA-3; instead, we replaced them with a more up-to-date sea-level dataset compiled by BOM, with a greater number of records. Next, we obtained measurements from 16 additional providers that were not in GESLA-2 (indicated by the grey shading in Table 1). GESLA-3 now includes higher-frequency sea-level data obtained from paper records via data archaeology (DA) exercises. These included 21 records in the United States collated by Bromirski et al. (2003), Talke et al. (2014, 2018, 2020, 2021), Familkhalili and Talke (2016), Chant et al. (2018), and Ray and Talke (2019), 5 records in the United Kingdom by Haigh et al. (2009) and 3 records in Spain digitized by Marcos et al. (2013, 2021). These datasets include the earliest higher-frequency data available for the Pacific Ocean (Astoria, 1855–1876 and San Francisco, 1858–1877) and stations on the US East Coast and Europe from the late 19th century. While some information such as the datum and time zone is available in GESLA-3 metadata for these DA sources, users are referred to the references above for more detailed discussions of data provenance and quality.

For five of the 36 sources within GESLA-3 (i.e., UHSLC, NOAA, NHS, MI-C, and MI-R), we downloaded the data automatically and rapidly via an Application Programming Interface (API). For the NHS dataset, we combined the more recent data from the late 1980's, downloaded via API, with historical data going back as far as 1915, that was provided to us directly. For 25 of the 36 sources, we manually downloaded the data from provider websites. For some providers, the data could be downloaded in bulk. However, for other providers, the data had to be downloaded one record at a time. Furthermore, for a few providers, the data had to be downloaded in 1–15-year blocks, for each record. For the remaining six sources (i.e., DA, DMI, NOC, ESEAS, ICG, and UZ), we obtained the data directly from the provider or copied the data from GESLA-2 (when updates were not available). The United States providers USGS, CDWR, SFWMD, NFWFMD, and NCDEM, and the Dutch provider RWS, did not discern between tidally influenced gauges and river-only gauges; in these cases, we hand-selected stations where there was the obvious presence of tidal forcing during at least part of the year and we did not include the river-only records. The

TABLE 2 Number of records and years of data for each data source

Number	Abbreviated name	No. of records	No. of years	No. records >100years	No. records 50–100years	No. records 20–50years	No. records 10–20years	No. records 5–10years	No. records <5years	Min year	Max year	No. of countries
1	UHSLC	692	17,843	17	86	228	136	104	121	1846	2019	97
2	NOAA	1,395	14,884	14	89	118	93	100	981	1897	2021	1
3	CMEMS	590	9,753	8	35	106	157	166	118	1886	2021	24
4	MEDS	868	8,761	7	63	57	49	63	629	1895	2021	1
5	USGS	464	5,314	0	0	23	263	118	60	1987	2021	1
6	BOM	125	4,603	2	26	85	11	1	0	1897	2020	14
7	RWS	124	4,482	4	17	54	27	12	10	1800	2018	1
8	JODC_JMA	81	3,475	0	52	15	12	2	0	1960	2019	1
9	SMHI	63	2,547	9	11	8	18	13	4	1851	2021	1
10	REFMAR	108	2,479	2	10	30	40	14	12	1821	2021	1
11	BODC	46	1,879	1	14	28	3	0	0	1915	2021	1
12	CDWR	98	1,877	0	0	40	31	20	7	1982	2021	1
13	JODC_JCG	31	1,667	0	22	9	0	0	0	1910	2019	2
14	NHS	24	1,503	2	15	6	1	0	0	1914	2020	1
15	JODC_GIAJ	25	1,402	0	15	10	0	0	0	1932	2019	1
16	WSV	66	1,262	0	0	44	10	11	1	1994	2020	1
17	JODC_PAHB	70	1,117	0	2	20	24	23	1	1961	2019	1
18	SFWMD	34	1,090	0	1	26	7	0	0	1970	2020	1
19	ISPRA	36	926	0	0	26	5	5	0	1971	2021	1
20	IEO	12	714	0	10	1	1	0	0	1943	2015	1
21	DA	29	685	1	3	10	3	3	9	1855	2019	3
22	UNAM	35	663	0	2	14	6	4	9	1946	2018	1
23	FMI	14	657	0	12	1	0	1	0	1971	2021	1
24	DMI	3	331	2	1	0	0	0	0	1891	2020	1
25	BFG	5	242	0	2	3	0	0	0	1917	2021	1
26	MILC	21	182	0	0	0	11	3	7	2006	2021	1
27	CCO	15	173	0	0	1	9	4	1	1996	2021	1
28	NOC	7	137	0	0	3	4	0	0	1957	2018	3
29	NWFWMD	9	100	0	0	0	5	4	0	2000	2021	1
30	ESEAS	5	63	0	0	1	1	0	3	1935	2003	3

(Continues)

TABLE 2 (Continued)

Number	Abbreviated name	No. of records	No. of years	No. records >100years	No. records 50–100years	No. records 20–50years	No. records 10–20years	No. records 5–10years	No. records <5years	Min year	Max year	No. of countries
31	CG	1	51	0	1	0	0	0	0	1970	2020	1
32	UZ	1	44	0	0	1	0	0	0	1974	2017	1
33	NCDEM	10	39	0	0	0	0	1	9	2013	2021	1
34	CV	1	38	0	0	1	0	0	0	1983	2020	1
35	MI_R	9	32	0	0	0	0	0	9	2018	2021	1
36	GLOSS	2	6	0	0	0	0	0	2	1997	1999	1
-	Total	5,119	91,021	69	489	969	927	672	1993	1800	2021	114

NOAA and MEDS datasets included records in the Great Lakes, and we retained these in GESLA-3.

In GESLA-1 and GESLA-2, we focused primarily on obtaining long records. However, many shorter records (a few days to a few years) are now being routinely provided by data centres. Furthermore, as described in Section 1, the GESLA dataset is increasingly being used for a wider range of analysis purposes. Short records, even those up to a month in duration, can be proved useful for a variety of applications, including the calculation of harmonic constituents and the validation of numerical models. Therefore, for GESLA-3, we included all the higher-frequency records that were available from the 36 providers, as long as they had at least 30 days of measurements. As discussed below, the inclusion of short records is a primary reason why the number of records and years greatly increased in GESLA-3, compared to GESLA-2.

For most sources we obtained the so-called “delayed mode” or “research quality” data, which typically becomes available to a user with a delay from days to years, enabling the data centres to perform quality control and include flags to highlight periods of good, suspect, and bad data values. The latest years available for each source are listed in Table 2. For around half of the sources, we obtained data up to October 2021 (the dates we did the final processing of the dataset). Most other datasets included data until the end of 2019 or 2020.

The number of records and years of data in GESLA-3 is listed in Table 2 for each of the 36 contributing sources. In total, GESLA-3 contains 91,021 years from 5,119 records. A map showing the locations of the records for GESLA-3 is shown in Figure 1. The areas where the coverage has most improved, compared to GESLA-2, are North America (Figure 2a), Europe (Figure 2b), Japan (Figure 2c), and Australia (Figure 2d). This is illustrated clearly in Figure 3, which shows the location of new records in GESLA-3 that are more than 50 km from a record in GESLA-2. Coverage outside of these regions is primarily achieved by ingesting the UHSLC dataset, which continues to be updated with new data, but has remained consistent in terms of the number and location of included stations. Coverage in North America has increased enormously for several reasons. First, we added all datasets available from NOAA and MEDS, not just the longer datasets. Furthermore, we also incorporated new datasets from the USGS, CDWR, SFWMD, NFWMD, NCDEM, and UNAM. In Europe, the largest increase in coverage stems from the records added from CMEMS. However, note many of the records from CMEMS only cover more recent decades, and not the full period often available from other providers (e.g., for Newlyn, data are available from 1915 from BODC, but only from 1990 from CMEMS). We also added new datasets for the United Kingdom from the

CCO, for Ireland from MI-R and MI-C, and for Germany from WSV. Coverage has increased significantly in Japan, from 80 records in GESLA-2 to 207 in GESLA-3. GESLA-2 only included data from the JODC_JMA. In GESLA-3, we have added data from the JODC_JCG, JODC_GIAJ, and JODC_PAHB. For Australia, the number of records has increased from 47 to 125, resulting in the development of the Australian National Collection of Homogenized Observations of Relative Sea Level (ANCHORS, Hague et al., 2021). The ANCHORS methodology applied statistical techniques to remove stepwise changes in annual means resulting, for example, from datum shifts and tide gauge relocations, for long tide-gauge records. So that quality control processes applied in GESLA-3 are internally consistent, we only included unhomogenized data from ANCHORS records, which is then quality controlled as described in Section 2.2. In the process of developing ANCHORS, many additional shorter records suitable for GESLA-3 were identified and are also included here.

In GESLA-3, records are available for 114 countries. The countries with the highest number of records are the United States and Canada, reflecting in part the vast length of the coastlines in these countries. The number of countries, covered by each of the 36 contributing sources is listed in Table 2 (final column). The UHSLC dataset contains records from 97 countries, significantly higher than any of the other sources. This illustrates how essential the UHSLC dataset is for achieving good global coverage in GESLA-3 (and earlier versions).

GESLA-3 contains 91,021 years of sea-level data (Table 2). The number of records containing different numbers of years is shown in Figure 4a. The record, with the most years of data (168 years between 1851 and 2021) is Olands Norra Udde from the SMHI, and the next longest record is Brest (165 years between 1846 and 2021) from REFMAR. The number of records with different ranges of years is shown in Figure 4b. The inclusion of many new short (i.e., <5 years) records is evident, but GESLA-3 also includes many new longer records, for example, for Japan from JODC_JCG, JODC_GIAJ, and JODC_PAHB, and for the United States and Europe from the DA sources. The record locations, with corresponding numbers of years, are shown in Figure 5. The majority of the sites with >100 years are located in North America and Europe. Four further sites are located in Panama and Australia. The number of records starting in particular year ranges is shown in Figure 4c. The location of records starting in the corresponding year ranges are shown in Figure 6. The earliest record, Katwijk in the Netherlands, starts in the year 1805 (but this record only contains 3 years). Hence, GESLA-3 spans the 217-year period from 1805 to 2021. The next earliest record, Saint Nazaire in France starts in the year 1821 (this record contains 134 years of data). The number of

records containing data each year between 1805 and 2021, is shown in Figure 4d, for GESLA-3, plotted alongside the same information for the earlier GESLA-1 and GESLA-2 datasets. There is an apparent decline in data availability in the most recent few years. As discussed above, this is because the “delayed mode” or “research quality” data become available with a typical delay of a few years during which data centres perform quality control.

2.2 | Data processing and format

The sea-level dataset we obtained from the 36 providers has differing units, time zones, and formats, and quality control flags are variously defined. As with GESLA-1 and GESLA-2, we converted height units to metres, the time zone of each record was adjusted to Coordinated Universal Time (UTC), we matched the specific data provider quality control flags to our defined GESLA flags (see below), and we processed the records into a standard format (a slightly modified version of the GESLA-2 format, see below). USGS and CDWR used Daylight Savings time in summer and we first shifted these to standard time, before converting to UTC; however, the times of annual shifts between Daylight Savings Time and Standard Time are imperfectly documented, some errors may remain.

In most instances, we did not adjust the frequency of the records, which in all cases was at least hourly, although several sources have data at higher frequency (6, 10, or 15 min). When given an option (e.g., on a provider's website), we always downloaded the hourly data, over higher-frequency data, as hourly data are adequate for most analyses that have previously been undertaken using GESLA, and it reduces the file sizes of the final processed datasets. Within the CMEMS dataset, the French data are provided at different frequencies for the same tide gauge. For example, the dataset at Brest is provided at 1-, 2-, 5-, 10-, and 60-min frequencies (for different overlapping periods). The higher-frequency records are generally much shorter, and the quality control is often less rigorous, so we ignored these and only included, in most instances, the hourly resolution dataset. The WSV data had a resolution of 1-min and the USGS, CDWR, SFWMD, NFWMD, and NCDEM data had resolutions between 1 and 15 min. We averaged these records, to hourly values, again to reduce the file size of the processed dataset. To do this, we selected all the data that lay within plus or minus 30 min from a specific hour, and averaged these values. Data from some providers are temporally regular (e.g., there is a date/time stamp every single hour) while for other providers the data are irregular (e.g., there is not a date/time stamp every hour – some are missing). In some cases, the frequency changes over time (e.g., the first part of the record




FIGURE 1 (a) Locations of the sea-level records in GESLA-3; with histograms of the record count plotted along (b) *y*-axis for latitude and (c) *x*-axis for longitude

is hourly, while the more recent period has a frequency of 15-min). We did not attempt to make the dataset temporally regular, or (with the exception of that mentioned above) adjust the frequency, as most analysis approaches can handle data with irregular time scales. Furthermore, we wanted the records to remain as consistent as possible with that provided by the originating agency.

For consistency, we have kept the format of the GESLA-3 data files virtually the same as in GESLA-2. As illustrated in Table 3, each text file contains 41 lines of header information, followed by the data itself. In GESLA-2, we listed only the name of the contributor of the data. However, in GESLA-3 we have included two extra header lines recording the website and the contact details of the contributor. For the international data centres, such as the UHSLC and CMEMS, the data they provide originate from different national centres. To ensure the originators of the data receive the credit they deserve, and so that the data can be traced back to the original providers, we have included three extra header lines listing the originator of the data, their website, and contact details. Where the contributor and originator are the same, the information is simply repeated. In GESLA-3, we have also added a new header line to indicate the record length in years.

We have also added a new header line to indicate the overall record quality, to aid the range of users of GESLA. A brief, qualitative expert judgement assessment was

made by visually inspecting every record in GESLA-3. Based on this evaluation, we now indicate if that record has (a) no obvious issues; (b) possible datum issues; (c) possible quality control issues; and (d) possible datum and quality control issues. In total, 4,747 records are classified as having no obvious issues, 149 as having possible datum issues, 179 as having possible quality control issues, and 46 as having possible datum and quality control issues. Users who want to assess trends in extreme sea levels might, for example, only use long records identified to have no obvious issues. By contrast, users who are interested in shorter time periods (e.g., for hydrodynamic model validation or investigation of a specific event) might choose to use all available records.

In GESLA-3, we have added many new records located in the upper reaches of estuaries and tidally influenced rivers, and we hope these new records may help spur scientific innovation in these dynamic, highly anthropogenically affected regions (see reviews by Hoitink and Jay, 2016; Haigh et al., 2020; and Talke and Jay, 2020). To aid in analysis, another new header line has therefore been added to indicate the hydrographic environment of the tide-gauge location. This header line denotes whether a record is associated with a (a) coastal, (b) river, or (c) lake environment. We visually inspected each record, and location, and distinguished between ‘coastal’ and ‘river’ stations based on whether the water level signal was clearly




FIGURE 2 Locations of the sea-level records in GESLA-2 and GESLA-3 for the four regions with the greatest coverage increase: (a) North America; (b) Europe; (c) Japan; and (d) Australia. Note the GESLA-2 locations are also in GESLA-3

dominated by tidal or river influences, considering the distance from the open coastline. “River” stations were classified as those where a strong river influence is evident in the water levels (and they are often some distance from the open coastline), whereas “coastal sites” were classified as those where the tidal component was clearly dominant. As mentioned earlier, if a record had no clear tidal signal, for at least part of the year, it was removed. Lake stations are in regions hydraulically disconnected from the ocean. The lake sites are mostly in the Great Lakes (from NOAA or MEDS), although a small selection of sites is in the IJsselmeer in the Netherlands (from RWS). We realize the subdivision into ‘coastal’ and ‘river’ is very difficult, and somewhat subjective, but we hope this is useful for users of the dataset. In total, 4,159 records are classified as coastal, 784 as a river, and 178 as a lake. Users only interested in assessing trends in extreme sea levels from oceanographic sources may wish to just select the coastal records, and ignore the records associated with river and lake stations.

In each file, the data itself is comprised of five columns, separated by one or more spaces, consistent with

GESLA-1 and GESLA-2. These are (a) the date, (b) time, (c) the observed sea level, (d) the quality control flag, and (e) the flag indicating whether the data should be used for analysis or not. Each data value in GESLA-3 has been assigned two flags. The first flag (in column 4) indicates the quality control undertaken by the provider. For this we use the following flags to be consistent with GESLA-1 and GESLA-2: 0 for no quality control; 1 for correct value; 2 for interpolated value; 3 for doubtful value; 4 for an isolated spike or wrong value, and 5 for missing value (set to -99.9999). Where available, we matched each of the provider flags to our system. Due to the huge effort it would require, we did not undertake further extensive quality control of our own. However, we did visually inspect each record individually, and we manually flagged suspect values that were clearly outside of the normal range or were isolated spikes. It is clear that data quality is poor for some sources, and datum jumps do exist, and users should treat these particular records with caution. As discussed earlier the overall record quality identifies the records that should be treated with caution. The second flag (in column 5) is a 1 or 0, indicating whether that value should




FIGURE 3 Location of sea-level records in GESLA-3 that are more than 50 km from a record in GESLA-2

be used for analysis, or not, respectively. All values whose quality control flag was 0, 1, or 2 were set to analysis flag 1 (use), and all values whose quality control flag was 3, 4, or 5, were set to analysis flag 0 (do not use).

The name of each file is made up of the (lower case) site name, site code, country code, and an abbreviation of the contributor name (note, for the DA records, we have added an underscore and the initials of the person who provided that record, for example, da_mm for the three records provided by Marta Marcos), separated by a hyphen (e.g., brest-822a-fra-uhsic). We have replaced all spaces in site names with an underscore. We have also removed all full stops, commas, brackets, accents, hyphens, and other special characters from file names and site codes. Hence, the file name and code might not exactly match that of the data provider. For country codes, we use the three-letter ISO 3166-1 alpha-3 codes (https://en.wikipedia.org/wiki/ISO_3166-1_alpha-3).

2.3 | Data licence


The developers of GESLA-1 only used data that had been provided to them on a personal basis, knowing how it was intended to be used. The dataset was subsequently made available only to trusted scientific users. For GESLA-2, the team divided the dataset into 27 “public” and 3 “private” sub-sets. Subject to the acknowledgement of the

data owner, the ‘public’ data set was readily available to download from the GESLA website and could be used for both research and consultancy purposes. However, the ‘private’ dataset could only be used for research and not a consultancy. This could only be obtained from the GESLA website with a password; bona fide researchers had to contact the GESLA team with an explanation of why they would like access to the dataset, to be given the password.

To simplify the process, we have decided not to separate the GESLA-3 data into two sets, on the GESLA website. Instead, we have examined the licences associated with each data contributor, where available, included a link to the specific licence in Table 1, and trust the users to comply with the licence conditions. Table 1 also lists whether the data can be freely used for research and/or consultancy. For example, users wishing to use the records provided by CV, UZ, and CMEMS for consultancy purposes, must contact these organizations to obtain permission first (or in the case of CMEMS the organizations that provided the data to them). In GESLA-2, the Australia records were included in the ‘private’ sub-set. However, we are pleased that in GESLA-3 permission has been obtained to make these Australian records publicly available.

In summary, the data are accessible but are covered by several different licences, some of which are noncommercial, by-attribution, or a combination of conditions. Access to the data does not currently require authentication, so restricted data are open to all, and we ask users to

FIGURE 4 (a) Number of records with the stated number of years of data in GESLA-3 (note the logarithmic scale); (b) the number of records with a particular number of years; (c) the number of records with data starting in a particular span of years; and (d) the number of records with data in a particular year in GESLA-3, -2 and -1 (note the logarithmic scale)



comply with the licence conditions. In acknowledgment of the central role of the UHSLC dataset in GESLA-3 (and earlier versions) and the decades-long effort to collect and quality assess the UHSLC data, we request that users of GESLA-3 data cite Caldwell et al. (2015) in addition to this paper in their work.

2.4 | Data principles

While constructing this third version of GESLA, we carefully considered the FAIR data principles, conceived by Wilkinson et al. (2016); that is that data should be findable, accessible, interoperable, and reusable. These principles also help ensure that proper credit is given to all those

involved in the data lifecycle. In GESLA-3, we have implemented several improvements compared to GESLA-2, to move the dataset towards being FAIR-compliant. The data archived with the BODC Published Data Library (PDL) has been assessed against the GO-FAIR criteria (<https://www.go-fair.org/fair-principles/>) and at the time of writing, partially meets the criteria. The GESLA-3 data are assigned a globally unique and persistent identifier and the metadata contains the identifier of the data (the DOI universally unique identifier, UUID, is given on the landing page). The datasets are findable in searchable resources, such as Google Dataset Search and included in metadata directories (e.g., the European Directory of Marine Environmental). The file header metadata have been improved since GESLA-2 as we have differentiated between




FIGURE 5 Locations of records with: (a) <5, station years; (b) 5–10; (c) 10–20; (d) 20–50; (e) 50–100; and (f) >100

who has contributed the data (# CONTRIBUTOR) and where the data originated from (# ORIGINATOR), but in the next version, we could look to implement more of the minimum mandatory metadata as detailed in the EuroSea deliverable D3.3 (Pérez Gómez et al., 2021).

We are working towards making the GESLA-3 data more interoperable. We have started to implement the use of some controlled vocabularies (e.g., ISO 3166-1 alpha-3 for country code), but in future versions we would like to include controlled vocabularies for other metadata. These would include using vocabularies such as the Research Organization Registry (<https://ror.org>) or the European Directory of Marine Organizations (<https://edmo.seadatanet.org>) for organizations, and SeaDataNet (<https://www.seadatanet.org>) for coordinate and datum information. The data can easily be converted into NetCDF (Network Common Data Form) files, and we hope to archive and distribute these data via an ERDDAP data server in the future, where allowable. We have also provided

computer scripts on the GESLA website in a variety of programming languages (e.g., MATLAB, Python, and R), to allow users to easily load in the dataset for scientific analysis.

3 | DATASET ACCESS

The 5,119 records in GESLA-3, and copies of the earlier two versions of the dataset, can be obtained from <https://www.gesla.org>. Furthermore, we now also provide a comma-delimited ASCII file containing information about each record and a Keyhole Markup Language (KML) file, which can be opened, for example, in Google Earth, to show record locations and information. On the GESLA website, we keep a list of any problems that we, or others, identify with the data, which we subsequently correct.

The GESLA-3 dataset has also been archived with the BODC, in two parts: the first part, which can be obtained




FIGURE 6 Locations of records with data starting in the years: (a) before 1850; (b) 1850–1900; (c) 1900–1950; (d) 1950–2000; (e) 2000–2015; and (f) after 2015

here https://www.bodc.ac.uk/data/published_data_library/catalogue/10.5285/d21a496a-a48e-1f21-e053-6c86abc08512/ contains the 4,527 records that can be used for both research and consultancy purposes and is covered by a creative commons CC-BY 4.0 licence and the second part, which can be downloaded here https://www.bodc.ac.uk/data/published_data_library/catalogue/10.5285/d21a496a-a48f-1f21-e053-6c86abc08512/ contains the 592 records that can be used for research purposes, but not consultancy and is covered by a creative commons BY-NC 4.0 licence.

4 | DISCUSSION AND CONCLUSIONS

This paper has described the assembly of the third version of the GESLA dataset. GESLA-3 is a major update,

containing 91,021 years of sea-level observations, more than double that of GESLA-2. The 5,119 records in GESLA-3 are nearly four times the number of that in GESLA-2. Many of the records are now available to October 2021, encompassing an extra 6 or 7 years of data compared to GESLA-2. Furthermore, new records have been added, improving spatial coverage, especially in North America, Europe, Japan, and Australia. In particular, we have added many new records for stations located in the upper reaches of estuaries and tidally influenced rivers.

There is some duplication between records provided by the different sources. For example, a record for Brest is provided by UHSLC, REFMAR, and CMEMS, and the data for Newlyn is provided by UHSLC, BODC, and CMEMS. Some duplicate records may be present in USGS and CDWR data, or NOAA and USGS. In some cases, two agencies may operate gauges within several kilometres

TABLE 3 Example of a GESLA-3 data file (brest-822a-france-uhslc) containing header lines followed by the hourly sea level values from UHSLC. A full description of the format is given at <https://www.gesla.org>

```
# FORMAT VERSION 5.0 Web: https://gesla.org Email: gesla.help@gmail.com
# SITE NAME Brest
# SITE CODE 822A
# COUNTRY FRA
# CONTRIBUTOR University of Hawaii Sea Level Center
# CONTRIBUTOR WEBSITE https://uhslc.soest.hawaii.edu
# CONTRIBUTOR CONTACT philiprt@hawaii.edu
# ORIGINATOR Systeme d'Observation du Niveau des Eaux Littorales (SONEL)
# ORIGINATOR WEBSITE Unspecified
# ORIGINATOR CONTACT Unspecified
# LATITUDE 48.38300000
# LONGITUDE -4.49500000
# COORDINATE SYSTEM Unspecified
# START DATE/TIME 1846/01/04 00:00:00
# END DATE/TIME 2018/12/31 23:00:00
# NUMBER OF YEARS 165
# TIME ZONE HOURS 0
# DATUM INFORMATION Unspecified
# INSTRUMENT Unspecified
# PRECISION Unspecified
# NULL VALUE -99.9999
# GAUGE TYPE Coastal
# OVERALL RECORD QUALITY No obvious issues
#
# CREATION DATE UTC 2021/11/01
#
# COLUMN 1 Date yyyy/mm/dd
# COLUMN 2 Time hh:mm:ss
# COLUMN 3 Observed sea level (m)
# COLUMN 4 Observed sea level QC flag
# COLUMN 5 Use-in-analysis flag (1 = use, 0 = do not use)
#
# Quality-control (QC) flags for column 4
#
# 0 - no quality control
# 1 - correct value
# 2 - interpolated value
# 3 - doubtful value
# 4 - isolated spike or wrong value
# 5 - missing value
#
1846/01/04 00:00:00      3.4800      1      1
1846/01/04 01:00:00      2.7000      1      1
1846/01/04 02:00:00      1.9900      1      1
1846/01/04 03:00:00      1.7000      1      1
<Followed by data to 2018/12/31 23:00:00>
```

of each other (e.g., the USGS and NOAA at Vancouver, Washington, or USGS and NOAA at Fort Pulaski, Georgia). The level of quality control may also differ between providers and the data lengths might not be consistent (e.g., the UHSLC and BODC dataset for Newlyn start in 1915 whereas the CMEMS record starts in 1990). At a tide gauge site with more than one record, we advise users to utilize the longest record, and preferably also the most up-to-date; a complementary strategy would be to use the agency giving the most attention to data quality (e.g., UHSLC in many cases) or the agency with the most experience measuring sea level (e.g., in a US context it is likely that NOAA has the most experience measuring sea level). Our choice to minimize data processing, and remain as consistent as possible with the originating agency, provides more freedom but also puts more responsibility on the end-user. We recommend, therefore, that researchers do due diligence and carry out additional quality assurance that is commensurate with their goals and needs. We are in the process of making a list of the tide gauge sites with duplicate records, and will make this available on the GESLA website in the future. We also hope to add derived products in the future (e.g., time series of astronomical tides and skew surges, etc.).

Despite the large improvement in the number of records and the number of years available, further improvements in the GESLA database are possible and desirable. As Woodworth et al. (2017) pointed out, GESLA-2 did not contain any data from India, for example, and there are only a few Bangladesh, Russian and Chinese sites made available via UHSLC. Mean sea-level data are available via PSMSL for these countries, but higher-frequency data are not distributed to the international community. A number of data series are only available commercially (e.g., from the National Mapping and Resource Information Authority [NAMRIA] in the Philippines or the Mekong Commission in Vietnam), and are therefore not included in GESLA-3. For example, only a fraction of the more than 1,000 years of data from the Philippines, spread over >50 stations, are available in GESLA-3 (through the UHSLC data set), though data can be purchased. Coverage across South America and Africa could also be better, although this primarily reflects a smaller number of operational stations rather than a lack of access to data. Additional records exist even in regions with high data coverage, for example for the Mississippi Delta from the US Army Corps of Engineers, or German authorities along the Ems River Estuary. Earlier digital records from our providers (Tables 1 and 2) are often unavailable online. For example, many USGS records from pre-2007 are unavailable (e.g., from Florida) due to uncertain data control. In Germany, many digital records are only in high water/low water format and are unavailable online; a similar issue exists

for data archaeology efforts (such as the high water/low water record from 1875 to the present made available in Ralston et al., 2019). In the future, the GESLA effort may therefore include a separate database for high water/low water or irregularly measured data, since these are often critical for assessing long-term trends in extremes (e.g., Dangendorf et al., 2013). Continued data archaeology efforts are needed; a number of records remain in nonelectronic format, even up to the 1980s, sometimes in formats only readable by specialized machines (e.g., Talke and Jay, 2017). Many thousands of years of additional records remain to be digitized, quality-assured, and published from around the Pacific Rim, North America, and Europe (e.g., Bradshaw et al., 2015; Pouvreau, 2008; Talke and Jay, 2013; Talke and Jay, 2017). Many historical records in other countries likely remain undocumented, undigitized, or otherwise unavailable. As these records become available, they will be added to the GESLA-3 database. Therefore, sea-level data archaeology efforts remain vital for improving 19th and 20th-century data coverage.

Due to the time-consuming nature of this work, updates to GESLA have been made in 5- or 6-year intervals. Because data providers have recently made it easier to obtain datasets via website downloads or APIs, we now hope to update the records more frequently. We also hope to continue to add new records from additional data providers, as we become aware of them. In GESLA-3, we have added, for the first time, 29 records captured recently from exercises in data archaeology; in the future, we hope to add many more records of this nature. We ask the readers and encourage data providers to contact us with details of any higher-frequency records that are available, but not currently in GESLA; we will endeavour to include these in future releases. As mentioned earlier, we also hope in the future to make GESLA data available via an ERDDAP data server.

While assembling GESLA-3, we became aware of a new sea-level dataset that has recently been assembled called MISELA (Minute Sea-Level Analysis) (Zemunik et al., 2021). This contains 1-minute sea-level data, at 331 tide gauges worldwide, required for studying oceanographic processes like seiches, meteotsunamis, infragravity, and coastal waves. We welcome this new dataset. Combined, the PSMSL, UHSLC, GESLA, and MISELA databases now allow for assessments of sea-level change across the full spectrum of frequencies of interest.

In concluding their paper, Woodworth et al. (2017) noted that the two scientists (Philip Woodworth and John Hunter), who provide the bulk of the construction of GESLA-2, had now retired. Now, under new leadership, the GESLA initiative continues, and the number of studies that use GESLA continues to grow. We are confident that further advances in the understanding of

ocean tides, storm surges, extreme sea levels, and other relevant coastal processes will stem from this new release and enhance insight into how coastal communities might respond to sea-level rise, extreme events, and climate change.

AUTHOR CONTRIBUTIONS

Ivan Haigh: Conceptualization (equal); data curation (equal); formal analysis (equal); methodology (equal); project administration (equal); visualization (equal); writing – original draft (equal); writing – review and editing (equal). **Marta Marcos:** Conceptualization (equal); data curation (supporting); formal analysis (supporting); writing – review and editing (supporting). **Stefan Andreas Talke:** Data curation (supporting); writing – review and editing (supporting). **Philip Woodworth:** Conceptualization (supporting); supervision (supporting); writing – review and editing (supporting). **John Hunter:** Conceptualization (supporting); supervision (supporting); writing – review and editing (supporting). **Ben Samuel Hague:** Data curation (supporting); writing – review and editing (supporting). **Arne Arns:** Data curation (supporting); writing – review and editing (supporting). **Elizabeth Bradshaw:** Project administration (supporting); writing – review and editing (supporting). **Philip Thompson:** Data curation (supporting); writing – review and editing (supporting).

ACKNOWLEDGEMENTS

We would like to wholeheartedly thank the data centres (listed in Table 1) and multiple individuals who provided the data in GESLA-3. We would specifically like to thank Andrew Matthews, Begoña Pérez Gómez, Marta de Alfonso Alonso-Muñoyerro, Angela Hibbert, Vibeke Huess, Ulpu Leijala, Guðmundur Birkir, Guy Westbrook, Torbjørn Taskjelle, Ruth Farre, Todd Ehret, Patrick Caldwell, Laurent Testut, Gerard McCarthy, Jenny Chiu, Chris Hughes, Ole Baltazar Andersen, Oleg Nikitin, Per Knudsen, Petra Zemunik, Thomas Dhoop, Charlie Thompson, and Scott Stephens (we are sorry if we have inadvertently missed anyone), for providing data or advice on various data related issues. The authors are grateful to Maritime Safety Queensland, Manly Hydraulics Laboratory, Victorian Regional Channels Authority, Western Australia Department of Transport, Victorian Ports Corporation Melbourne, Tasmanian Ports Corporation, Pilbara Ports Authority, Gippsland Ports, HydroSurvey (Flinders Ports), and Fremantle Ports for providing ongoing raw tide gauge data to the Bureau of Meteorology, which provides the basis for the quality-controlled hourly-resolution Australian contribution to GESLA-3.

We received no direct funding to assemble GESLA 3, however part of our time was funded on relevant grants,

as follows: IDH time was partly funded via the NERC-funded CHANCE Project (NE/S010262/1); SAT was partly funded by the National Science Foundation (Award number 1455350 and 2013280); MM was supported by European Regional Development Fund/Ministerio de Ciencia, Innovación y Universidades – Agencia Estatal de Investigación through the MOCCA project (grant no. RTI2018-093941-B-C31); PRT was supported by the National Oceanic and Atmospheric Administration Global Ocean Monitoring and Observation Program via the University of Hawai'i Sea Level Center (grant no. NA11NMF4320128).

CONFLICT OF INTEREST

The authors declare no conflicts of interest.

OPEN RESEARCH STATEMENT

This article has earned an Open Data badge for making publicly available the digitally-shareable data necessary to reproduce the reported results. The data is available at https://www.bodc.ac.uk/data/published_data_library/catalogue/10.5285/d21a496a-a48e-1f21-e053-6c86abc08512/ and https://www.bodc.ac.uk/data/published_data_library/catalogue/10.5285/d21a496a-a48f-1f21-e053-6c86abc08512/. Learn more about the Open Practices badges from the Center for Open Science: <https://osf.io/tvyxz/wiki>

ORCID

Ivan D. Haigh  <https://orcid.org/0000-0002-9722-3061>

Marta Marcos  <https://orcid.org/0000-0001-9975-5013>

Philip L. Woodworth  <https://orcid.org/0000-0002-6681-239X>

[org/0000-0002-6681-239X](https://orcid.org/0000-0002-6681-239X)

Ben S. Hague  <https://orcid.org/0000-0002-4931-8111>

REFERENCES

- Arns, A., Wahl, T., Wolff, C., Vafeidis, A.T., Haigh, I.D., Woodworth, P. et al. (2020) Non-linear interaction modulates global extreme sea levels, coastal flood exposure, and impacts. *Nature Communications*, 11, 1918. <https://doi.org/10.1038/s41467-020-15752-5>
- Bradshaw, L., Rickards, L. & Aarup, T. (2015) Sea level data archaeology and the Global Sea Level Observing System (GLOSS). *GeoResJ*, 6, 916. <https://doi.org/10.1016/j.grj.2015.02.005>
- Bromirski, P.D., Flick, R.E. & Cayan, D.R. (2003) Storminess variability along the California coast: 1858–2000. *Journal of Climate*, 16(6), 982–993. [https://doi.org/10.1175/1520-0442\(2003\)016<0982:SVATCC>2.0.CO;2](https://doi.org/10.1175/1520-0442(2003)016<0982:SVATCC>2.0.CO;2)
- Caldwell, P.C., Merrifield, M.A. Thompson, P.R. (2015), *Sea level measured by tide gauges from global oceans—the Joint Archive for Sea Level Holdings (NCEI Accession 0019568), Version 5.5, NOAA National Centers for Environmental Information, Dataset.* <https://doi.org/10.7289/V5V40S7W>
- Chant, R.J., Sommerfield, C.K. & Talke, S.A. (2018) Impact of channel deepening on tidal and gravitational circulation in a highly engineered estuarine basin. *Estuaries and Coasts*, 41(6), 1587–1600. <https://doi.org/10.1007/s12237-018-0379-6>
- Church, J.A., Clark, P.U., Cazenave, A., Gregory, J.M., Jevrejeva, S., Levermann, A. et al. (2013) Sea level change. In: Stocker, T.F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S.K., Boschung, J. et al. (Eds.) *Climate change 2013: The physical science basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge and New York, NY: Cambridge University Press. pp. 1137–1216.
- Dangendorf, S., Mudersbach, C., Wahl, T. & Jensen, J. (2013) Characteristics of intra-, inter-annual and decadal sea-level variability and the role of meteorological forcing: the long record of Cuxhaven. *Ocean Dynamics*, 63, 209–224.
- Familkhalili, R. & Talke, S.A. (2016) The effect of channel deepening on storm surge: A case study of Wilmington, NC. *Geophysical Research Letters*, 43(17), 9138–9147. <https://doi.org/10.1002/2016GL069494>
- Fox-Kemper, B., Hewitt, H.T., Xiao, C., Aðalgeirsdóttir, G., Drijfhout, S.S., Edwards, T.L. et al. (2021) Ocean, cryosphere and sea level change. In: Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S.L., Péan, C., Berger, S. et al. (Eds.) *Climate change 2021: The physical science basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge and New York, NY: Cambridge University Press. pp. 1211–1362. <https://doi.org/10.1017/9781009157896.011>
- Hague, B.S., Jones, D.A., Trewin, B., Jakob, D., Murphy, B., Martin, D. et al. (2021) ANCHORS: A multi-decadal tide gauge data sets to monitor Australian relative sea level changes. *Geoscience Data Journal*, 1–17. <https://doi.org/10.1002/gdj3.136>
- Haigh, I., Nicholls, R. & Wellsb, N. (2009) Mean sea level trends around the English Channel over the 20th century and their wider context. *Continental Shelf Research*, 29, 2083–2098.
- Haigh, I.D., Pickering, M.D., Green, J.A.M., Arbic, B.K., Arns, A., Dangendorf, S. et al. (2020) The tides they are a' changing: A comprehensive review of past and future non-astronomical changes in tides, their driving mechanisms and future implications. *Reviews of Geophysics*, 58(1), e2018RG000636. <https://doi.org/10.1029/2018RG000636>
- Hoitink, A.J.F. & Jay, D.A. (2016) Tidal river dynamics: Implications for deltas. *Reviews of Geophysics*, 54, 240–272. <https://doi.org/10.1002/2015RG000507>
- Holgate, S.J., Matthews, A., Woodworth, P.L., Rickards, L.J., Tamisiea, M.E., Bradshaw, E. et al. (2013) New data systems and products at the permanent service for mean sea level. *Journal of Coastal Research*, 29(3), 493–504. <https://doi.org/10.2112/JCOASTRES-D-12-00175.1>
- Hunter, J.R., Church, J.A., White, N.J. & Zhang, X. (2013) Towards a global regionally varying allowance for sea-level rise. *Ocean Engineering*, 71, 17–27. <https://doi.org/10.1016/j.oceaneng.2012.12.041>
- Hunter, J.R., Woodworth, P.L., Wahl, T. & Nicolls, R.J. (2017) Using global tide gauge data to validate and improve the representation of extreme sea levels in flood impact studies. *Global and Planetary Change*, 156, 34–45. <https://doi.org/10.1016/j.gloplacha.2017.06.007>
- Marcos, M. & Woodworth, P.L. (2017) Spatio-temporal changes in extreme sea levels along the coasts of the North Atlantic and the Gulf of Mexico. *Journal of Geophysical Research Oceans*, 122, 7031–7048. <https://doi.org/10.1002/2017JC013065>

- Marcos, M., Puyol, B., Calafat, F.M. & Woppelmann, G. (2013) Sea level changes at Tenerife Island (NE Tropical Atlantic) since 1927. *Journal of Geophysical Research, Oceans*, 118, 4899–4910. <https://doi.org/10.1002/jgrc.20377>
- Marcos, M., Calafat, F.M., Berihuete, Á. & Dangendorf, S. (2015) Long-term variations in global sea level extremes. *Journal of Geophysical Research, Oceans*, 120, 8115–8134. <https://doi.org/10.1002/2015JC011173>
- Marcos, M., Puyol, B., Amores, A., Pérez Gómez, B., Fraile, M.Á. & Talke, S.A. (2021) Historical tide gauge sea-level observations in Alicante and Santander (Spain) since the 19th century. *Geoscience Data Journal*, 8, 144–153. <https://doi.org/10.1002/gdj3.112>
- Mawdsley, R.J., Haigh, I.D. & Wells, N.C. (2015) Global secular changes in different tidal high water, low water and range levels. *Earth's Future*, 3, 66–81. <https://doi.org/10.1002/2014EF000282>
- Menendez, M. & Woodworth, P.L. (2010) Changes in extreme high water levels based on a quasi-global tide-gauge dataset. *Journal of Geophysical Research*, 115, C10011. <https://doi.org/10.1029/2009JC005997>
- Muis, S., Irazoqui Apecechea, M., Dullaart, J., de Lima Rego, J., Madsen, K.S., Su, J. et al. (2020) A high-resolution global dataset of extreme sea levels, tides, and storm surges, including future projections. *Frontiers in Marine Science*, 7, 263. <https://doi.org/10.3389/fmars.2020.00263>
- Oppenheimer, M., Glavovic, B.C., Hinkel, J., van de Wal, R., Magnan, A.K., Abd-Elgawad, A. et al. (2019) Sea level rise and implications for low lying islands, coasts and communities. In: Pörtner, H.O., Roberts, D.C., MassonDelmotte, V., Zhai, P., Tignor, M., Poloczanska, E. et al. (Eds.) *IPCC Special Report on the Ocean and Cryosphere in a Changing Climate*. Cambridge and New York, NY: Cambridge University Press. pp. 321–445. <https://doi.org/10.1017/9781009157964.006>
- Peng, D., Hill, E.M., Meltzner, A.J. & Switzer, A.D. (2019) Tide gauge records show that the 18.61-year nodal tidal cycle can change high water levels by up to 30 cm. *Journal of Geophysical Research: Oceans*, 124(1), 736–749. <https://doi.org/10.1029/2018JC014695>
- Pérez Gómez, B., Testut, L., Hibbert, A., Matthews, A., Bradshaw, E., Westbrook, G., et al., (2021). *EuroSea Deliverable D3.3: New Tide Gauge Data Flow Strategy*. https://eurosea.eu/download/outputs_and_reports/deliverables/EuroSea-D3.3_New_Tide_Gauge_Data_Flow_Strategy.pdf
- Piccioni, G., Dettmering, D., Passaro, M., Schwatke, C., Bosch, W. & Seitz, F. (2018) Coastal improvements for tide models: the impact of ALES retracker. *Remote Sensing*, 10, 700. <https://doi.org/10.3390/rs10050700>
- Piccioni, G., Dettmering, D., Bosch, W. & Seitz, F. (2019) TICON: Tidal Constants based on GESLA sea-level records from globally located tide gauges. *Geoscience Data Journal*, 6, 97–104. <https://doi.org/10.1002/gdj3.72>
- Pouvreau, N., (2008). *Trois cents ans de mesures marégraphiques en France: outils, méthodes et tendances des composantes du niveau de la mer au port de Brest*. Ph.D. thesis, Université de La Rochelle.
- Ralston, D.K., Talke, S.A., Geyer, W.R., Al'Zubadaei, H. & Sommerfield, C.K. (2019) Bigger tides, less flooding: Effects of dredging on water level in the Hudson River estuary. *Journal of Geophysical Research*, 124(1), 196–211. <https://doi.org/10.1029/2018JC014313>
- Rashid, M.M., Wahl, T. & Chambers, D.P. (2021) Extreme sea level variability dominates coastal flood risk changes at decadal time scales. *Environmental Research Letters*, 16, 24026.
- Ray, R.D. (2020) First global observations of third-degree ocean tides. *Science Advances*, 6, eabd4744.
- Ray, R. & Talke, S.A. (2019) Nineteenth-Century Tides in the Gulf of Maine and implications for secular trends. *Journal of Geophysical Research*, 124, 7046–7067. <https://doi.org/10.1029/2019JC015277>
- Rhein, M., Rintoul, S.R., Aoki, S., Campos, E., Chambers, D., Feely, R.A. et al. (2013) Observations: Ocean. In: Stocker, T.F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S.K., Boschung, J. et al. (Eds.) *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge, New York, NY: Cambridge University Press.
- Santamaria-Aguilar, S. & Vafeidis, A.T. (2019) Are extreme skew surges independent of high water levels in a mixed semidiurnal tidal regime? *Journal of Geophysical Research*, 123, 8877–8886. <https://doi.org/10.1029/2018JC014282>
- Schindelegger, M., Green, J.A.M., Wilmes, S.-B. & Haigh, I.D. (2018) Can we model the effect of observed sea level rise on tides? *Journal of Geophysical Research Oceans*, 123, 4593–4609. <https://doi.org/10.1029/2018JC013959>
- Shalowitz, A.L. (1962) *Shore and sea boundaries: With special reference to the interpretation and use of coast and geodetic survey data* (Vol. 1). Washington, DC: Government Printing Office.
- Tadesse, M., Wahl, T. & Cid, A. (2020) Data-driven modeling of global storm surges. *Frontiers in Marine Science*, 7, 260. <https://doi.org/10.3389/fmars.2020.00260>
- Talke, S.A. & Jay, D.A. (2013) Nineteenth Century North American and Pacific Tides: Lost or just forgotten? *Journal of Coastal Research*, 29(6a), 118–127.
- Talke, S.A., D.A. Jay (2017). *Archival water-level measurements: recovering historical data to help design for the future*. US Army Corps of Engineers: Civil Works Technical Series, Report CWTS-02, 49p.
- Talke, S.A. & Jay, D.A. (2020) Changing tides: The role of natural and anthropogenic factors. *Annual Review of Marine Science*, 12, 121–151. <https://doi.org/10.1146/annurev-marine-010419-010727>
- Talke, S.A., Orton, P. & Jay, D.A. (2014) Increasing storm tides in New York Harbor, 1844–2013. *Geophysical Research Letters*, 41(9), 3149–3155. <https://doi.org/10.1002/2014GL059574>
- Talke, S.A., Kemp, A. & Woodruff, J. (2018) Relative sea level, tides, and extreme water levels in Boston (MA) from 1825 to 2018. *Journal of Geophysical Research*, 123(6), 3895–3914. <https://doi.org/10.1029/2017JC013645>
- Talke, S.A., Mahedy, A., Jay, D.A., Lau, P., Hillely, C. & Hudson, A. (2020) Sea level, tidal and river flow trends in the Lower Columbia River Estuary, 1853–present. *Journal of Geophysical Research-Oceans*, 125, 1–29. <https://doi.org/10.1029/2019JC015656>
- Talke, S.A., Familkhalili, R. & Jay, D.A. (2021) The influence of channel deepening on tides, river discharge effects, and storm surge. *Journal of Geophysical Research: Oceans*, 126, 1–24. <https://doi.org/10.1029/2020JC016328>

- Tsitsikas, C. (2018). *Regional sea level allowances along the world coast-line*. Master's Thesis, Utrecht, Netherlands, Utrecht University.
- Wahl, T., Haigh, I.D., Nicholls, R.J., Arns, A., Dangendorf, S., Hinkel, J. et al. (2017) Understanding extreme sea levels for broad-scale coastal impact and adaptation analysis. *Nature Communications*, 8, 1–12.
- Ward, P.J., Couasnon, A., Eilander, D., Haigh, I.D., Hendry, A., Muis, S. et al. (2018) Dependence between high sea-level and high river discharge increases flood hazard in global deltas and estuaries. *Environmental Research Letters*, 13, 84012. <https://doi.org/10.1088/1748-9326/aad400>
- Wilkinson, M., Dumontier, M., Aalbersberg, I., Appleton, G., Axton, M., Baak, A. et al. (2016) The FAIR Guiding Principles for scientific data management and stewardship. *Scientific Data*, 3, 160018. <https://doi.org/10.1038/sdata.2016.18>
- Wolff, C., Vafeidis, A.T., Muis, S., Lincke, D., Satta, A., Lionello, P. et al. (2018) A Mediterranean coastal database for assessing the impacts of sea-level rise and associated hazards. *Scientific Data*, 5, 180044. <https://doi.org/10.1038/sdata.2018.44>
- Wong, P.P., Losada, I.J., Gattuso, J.-P., Hinkel, J., Khattabi, A., McInnes, K.L. et al. (2014) Coastal systems and low-lying areas. In: Field, C.B., Barros, V.R., Dokken, D.J., Mach, K.J., Mastrandrea, M.D., Bilir, T.E. et al. (Eds.) *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge and New York, NY: Cambridge University Press, pp. 361–409.
- Woodworth, P.L. & Hibbert, A. (2018) The nodal dependence of long-period ocean tides in the Drake Passage. *Ocean Science*, 14, 711–730. <https://doi.org/10.5194/os-14-711-2018>
- Woodworth, P.L., Hunter, J.R., Marcos, M., Caldwell, P., Menendez, M. & Haigh, I.D. (2017) Towards a global higher-frequency sea level data set. *Geoscience Data Journal*, 3(2), 50–59. <https://doi.org/10.1002/gdj3.42>
- Woodworth, P.L., Melet, A., Marcos, M., Ray, R.D., Wöppelmann, G., Sasaki, Y.N. et al. (2019) Forcing factors affecting sea level changes at the coast. *Surveys in Geophysics*, 40, 1351–1397.
- Woodworth, P.L., Hunter, J.R., Marcos, M. & Hughes, C.W. (2021) Towards reliable global allowances for sea level rise. *Global and Planetary Change*, 203, 103522. <https://doi.org/10.1016/j.gloplacha.2021.103522>
- Zemunik, P., Šepić, J., Pellikka, H., Čatipović, L. & Vilibić, I. (2021) Minute Sea-Level Analysis (MISELA): A high-frequency sea-level analysis global dataset. *Earth System Science Data (ESSD)*, 13, 4121–4132. <https://doi.org/10.5194/essd-13-4121-2021>

How to cite this article: Haigh, I.D., Marcos, M., Talke, S.A., Woodworth, P.L., Hunter, J.R. & Hague, B.S. et al. (2022) GESLA Version 3: A major update to the global higher-frequency sea-level dataset. *Geoscience Data Journal*, 00, 1–22. Available from: <https://doi.org/10.1002/gdj3.174>